

11-6-2015

Peroxide Content of Secondary Organic Aerosol

Matthew E. Wise

Concordia University - Portland, mawise@cu-portland.edu

Ryan Caylor

Concordia University - Portland

Follow this and additional works at: <http://commons.cu-portland.edu/mathscienceresearch>



Part of the [Environmental Chemistry Commons](#)

Recommended Citation

Wise, Matthew E. and Caylor, Ryan, "Peroxide Content of Secondary Organic Aerosol" (2015). *Undergraduate Research*. 4.
<http://commons.cu-portland.edu/mathscienceresearch/4>

This Poster is brought to you for free and open access by the Math & Science Department at CU Commons. It has been accepted for inclusion in Undergraduate Research by an authorized administrator of CU Commons. For more information, please contact libraryadmin@cu-portland.edu.

Concordia University, Portland Oregon

From the SelectedWorks of Matthew E. Wise

November 6, 2015

Peroxide Content of Secondary Organic Aerosol

Ryan Caylor
Matthew E. Wise
John Shilling



SELECTEDWORKS™

Available at: http://works.bepress.com/concordia_wise_group/10/



Abstract

Secondary organic aerosol (SOA) formed from the oxidation of monoterpenes can impact the Earth's radiation balance, act as cloud condensation nuclei and negatively affect human health. In the summer of 2015, the **Secondary Organic Aerosol From Forest Emission Experiment (SOAFFEE)** laboratory campaign was launched in order to study the physical properties of SOA generated from the oxidation of α -pinene and Δ -carene. Both compounds are gas-phase monoterpenes emitted into the atmosphere via biogenic sources. In this study, the peroxide content of SOA was determined using an iodometric-spectrophotometric (IS) technique. It was found that the peroxide content of SOA generated during the SOAFFEE campaign was similar to that found in previous studies.

Possible SOA Reaction Mechanisms




Figure 1. Possible SOA reaction mechanisms (adapted from Docherty et al., 2005).

Environmental Chamber




Figure 2. Schematic of the environmental chamber

Materials and Methods





Figure 3. Reactions used in the IS technique. Colorless H_2O_2 is converted into yellow colored I_3^- ions. The absorbance of the I_3^- ions is measured using a high resolution UV/Vis spectrometer.




Figure 4. Experimental set-up used for the IS reactions.




Figure 5. UV-Vis absorption spectrum of 15.62 μM H_2O_2 and SOA. The SOA was created using 10 ppb α -pinene, 1 ppm H_2O_2 and 5 ppb NO .

Results




Figure 6. Hydrogen peroxide calibration curve ($A_{470} - A_{542}$)

VOC	VOC (ppb)	H2O2 (ppm)	O3 (ppb)	NO (ppb)	Cyclohexane/MeOH (ppm)	Peroxide content (wt%), 300 g/mol
a-pinene	10	0	80	0	0	18.85
a-pinene	10	0	80	0	10	89.17
a-pinene	10	0	80	25	10	21.09
a-pinene	10	0	80	25	10	15.74
a-pinene	10	1	0	0	0	47.70
a-pinene	10	1	0	1	0	51.78
a-pinene	10	1	0	5	0	26.00
a-pinene	10	1	0	10	0	25.66
a-pinene	10	1	0	25	0	5.80
a-pinene	10	1	60	0	0	46.95
a-pinene	20	1	60	0	0	18.50
a-pinene	20	1	60	0	0	16.16
a-pinene	20	1	60	0	0	43.92
a-pinene	20	1	60	0	0	25.25
a-pinene	20	1	60	0	0	36.72
a-pinene	40	2	60	0	0	64.93
d-carene	10	0.25	0	0	0	47.49
d-carene	10	0.5	0	0	0	41.67
d-carene	10	1	0	0	0	26.24

Table 1. Estimated peroxide content in SOA generated during the SOAFFEE campaign.

Comparison With Literature

monoterpene	OH scavenger	RH (%)	$Y_{\text{SOA/MT}}^a$ (%)	$Y_{\text{OP/MT}}^b$ (%)	$Y_{\text{OP/SOA}}^c$ (%)
α -pinene	cyclohexane	<0.5	48.7	4.6	28.6
		<0.5	45.2	7.5	49.8
		<0.5	50.6	5.2	30.8
		<0.5	39.5	7.3	55.5
		~50	51.9	7.1	41.3
1-propanol	cyclohexane	<0.5	41.5	7.9	57.3
		<0.5	48.8	7.0	43.0
		<0.5	47.1	10.1	64.5
formaldehyde	cyclohexane	<0.5	42.4	6.0	42.7
		<0.5	29.5	6.0	60.8
Δ - ³ carene	cyclohexane	<0.5	47.5	3.9	24.5
		<0.5	55.4	8.0	43.5
β -pinene	cyclohexane	<0.5	32.5	8.6	79.2
		<0.5	31.7	7.9	74.7
		~50	21.1	5.1	72.8
		~50	21.6	6.7	93.6
		<0.5	15.1	6.0	119.9
1-propanol	cyclohexane	<0.5	12.4	3.2	76.7
		<0.5	10.2	3.0	88.3
		<0.5	8.3	2.1	77.1
sabinene	cyclohexane	<0.5	38.1	11.7	92.6
		<0.5	10.4	3.6	103.1

^a Mass yield of SOA relative to reacted monoterpene (SOA (mass)/reacted monoterpene (mass)) $\times 100$. ^b Molar yield of SOA organic peroxides relative to reacted monoterpene (SOA peroxide (mol)/reacted monoterpene (mol)) $\times 100$. ^c Mass yield of SOA organic peroxides relative to SOA [(SOA peroxide (mol) \times MW (300 g mol⁻¹))/SOA mass (g)] $\times 100$.

Table 2. Measured SOA and organic peroxide yields from reactions of monoterpenes with ozone (Docherty et al., 2005).

Conclusions

The IS technique, used to measure the total peroxide content of SOA, was optimized. The hydrogen peroxide calibration curve constructed during the optimization process agreed well with the literature. SOA created during the SOAFFEE study were analyzed to determine their peroxide content. It was found that the SOA particles had similar amounts of peroxide as those created in previous experimental studies. The results presented here will aid in the determination of SOA composition and formation pathways. However, it is not sufficient to fully quantify the peroxide content of SOA. More UV/Vis absorbance measurements must be collected using SOA generated under different conditions. Furthermore, it is not clear if a hydrogen peroxide calibration curve is sufficient to predict total peroxide content in all types of SOA. Additional experimental methods can be used to validate the IS procedure.

References

- Mertes, P., Pfaffenberger, L., Dommen, J., Kalberer, M., Baltensperger, U. (2012). Development of a sensitive long path absorption photometer to quantify peroxides in aerosol particles (Peroxide-LOPAP). *Atmospheric Measurement Techniques*, 5, 2339-2348.
- Docherty, K.S., Wu, W., Lim, Y.B., and Ziemann, P.J. (2005). Contributions of organic peroxides to secondary aerosol formed from reactions of monoterpenes with O_3 . *Environmental Science and Technology*, 39, 4049-4059.
- Xu, L., Kollman, M. S., Song, C., Shilling, J. E., & Ng, N. L. (2014). Effects of NO_x on the volatility of secondary organic aerosol from isoprene photooxidation. *Environmental Science & Technology*, 48(4), 2253-2262.

Acknowledgements

- Visiting Faculty Program at the Pacific Northwest National Laboratory
- Concordia University Summer Undergraduate Research Institute